

Output results of CLIME (CLustering by Inferred Models of Evolution)

Dataset:

Num of genes in input gene set: 5
Total number of genes: 20834
Prediction LLR threshold: 0

The CLIME PDF output two sections:


1) Overview of Evolutionarily Conserved Modules (ECMs)

- Top panel shows the predefined species tree.
- Bottom panel shows the partition of input genes into Evolutionary Conserved Modules (ECMs), ordered by ECM strength (shown at right), and separated by horizontal lines.
- Each row show one gene, where the phylogenetic profile indicates presence (blue) or absence (gray) of homologs in each species (column).
- Gene symbols are shown at left. Gray color indicates that the gene is a paralog to a higher scoring gene within the same ECM (based on BLASTP $E < 1e-3$).


2) Details of each ECM and its expansion ECM+

- Top panel shows the inferred evolutionary history on the predefined species tree. Branch color shows the gain event (blue) and loss events (red color, with brighter color indicating higher confidence in loss). Branches before the gain or after a loss are shown in gray.
- Bottom panel shows the input genes that are within the ECM (blue/white rows) as well as all genes in the expanded ECM+ (green/gray rows). The ECM+ includes genes likely to have arisen under the inferred model of evolution relative to a background model, and scored using a log likelihood ratio (LLR).
- PG indicates "paralog group" and are labeled alphabetically (i.e., A, B). The first gene within each paralog group is shown in black color. All other genes sharing sequence similarity (BLAST $E < 1e-3$) are assigned to the same PG label and displayed in gray.

Overview of Evolutionarily Conserved Modules (ECMs)




Num of ECM Genes: 2. Num of Predicted Genes: 100. ECM Strength: 5.9




1: Valine, leucine and isoleucine degradation || 2: Fatty acid elongation || 3: Fatty acid metabolism || 4: Biosynthesis of unsaturated fatty acids || 5: alpha-Linolenic acid metabolism || 6: Steroid hormone biosynthesis || 7: Sulfur metabolism || 8: Pyruvate metabolism || 9: Linoleic acid metabolism || 10: Fructose and mannose metabolism || 11: Butanoate metabolism || 12: Ribosome biogenesis in eukaryotes || 13: Terpenoid backbone biosynthesis || 14: Phagosome || 15: Arginine and proline metabolism || 16: Glycolysis / Gluconeogenesis || 17: Glycerolipid metabolism

Num of ECM Genes: 2. Num of Predicted Genes: 100. ECM Strength: 5.9



3 / 4
4
3 / 4
5


Num of ECM Genes: 2. Num of Predicted Genes: 100. ECM Strength: 5.9



1: Sphingolipid metabolism || 2: Valine, leucine and isoleucine degradation || 3: Terpenoid backbone biosynthesis || 4: Purine metabolism || 5: Pyrimidine metabolism || 6: RNA degradation || 7: Apoptosis

ECM 2, Gene set "Fatty acid elongation", Page 1



Num of ECM Genes: 2. Num of Predicted Genes: 0. ECM Strength: 3.7

[illegible]

1: Lysosome || 2: Fatty acid elongation

ECM 3, Gene set "Fatty acid elongation", Page 1


Num of ECM Genes: 1. Num of Predicted Genes: 100



1: Fatty acid elongation || 2: Glycerophospholipid metabolism || 3: Glyoxylate and dicarboxylate metabolism || 4: Propanoate metabolism || 5: Valine, leucine and isoleucine degradation || 6: Arachidonic acid metabolism || 7: Linoleic acid metabolism ||
8: Steroid biosynthesis || 9: Pyruvate metabolism || 10: Fatty acid biosynthesis || 11: p53 signaling pathway || 12: Pyrimidine metabolism

ECM 3, Gene set "Fatty acid elongation", Page 2

Num of ECM Genes: 1. Num of Predicted Genes: 100

[illegible]

1: Primary bile acid biosynthesis || 2: Retinol metabolism || 3: Arachidonic acid metabolism || 4: Fatty acid metabolism || 5: Steroid biosynthesis || 6: Steroid hormone biosynthesis

Num of ECM Genes: 1. Num of Predicted Genes: 100

